

## **Euscorpius sicanus (Scorpiones: Euscorpiidae) from Tunisia: DNA barcoding confirms ancient disjunctions across the Mediterranean Sea**

Matthew R. Graham<sup>1</sup>, Pavel Stoev<sup>2</sup>, Nesrine Akkari<sup>3</sup>,  
Gergin Blagoev<sup>4</sup> & Victor Fet<sup>5</sup>

<sup>1</sup> School of Life Sciences, University of Nevada, Las Vegas, 4505 South Maryland Parkway,  
Nevada 89154-4004, USA

<sup>2</sup> National Museum of Natural History, Tsar Osvoboditel Blvd. 1, Sofia 1000, Bulgaria

<sup>3</sup> Natural History Museum of Denmark, University of Copenhagen, Universitetsparken 15,  
DK-2100 København Ø – Denmark

<sup>4</sup> Biodiversity Institute of Ontario, University of Guelph, Guelph, Ontario N1G 2W1, Canada

<sup>5</sup> Department of Biological Sciences, Marshall University, Huntington,  
West Virginia 25755-2510, USA

### **Abstract**

We used a DNA barcoding marker (mitochondrial *cox1*) to investigate the controversial natural occurrence of *Euscorpius sicanus* (C.L. Koch) in North Africa. We tested this hypothesis by comparing a sample collected from a mountain in Tunisia to disjunct populations in Sardinia, Malta, and Greece. Using these samples, and a few additional *Euscorpius* spp. from southern Europe as outgroups, we reconstructed the maternal phylogeny. We then used a molecular clock to place the phylogeny in a temporal context. The Tunisian sample grouped closest to a specimen from Sardinia, with both being more distantly related to *E. sicanus* from Malta, which is known to be genetically similar to samples from Sicily. Molecular clock estimates suggest an ancient disjunction across the Mediterranean Sea, with the divergence between samples from Sardinia and Tunisia estimated to have occurred between the Late Miocene and late Pliocene. The divergence date (mean = 5.56 Mya) closely corresponds with the timing of a sudden refilling of the Mediterranean Sea after it had evaporated during the Messinian salinity crisis. This rapid influx of water, in conjunction with tectonic activity, could have sundered connections between *Euscorpius* in North Africa and what is now the island of Sardinia. These results provide yet another case in which DNA barcodes have proven useful for more than just identifying and discovering species.

**Keywords:** Zanclean Flood, post-Messinian Flood, Messinian salinity crisis, molecular clock, *cox1*, mitochondrial DNA, barcode.

## Introduction

Most of North Africa's rich scorpion fauna, which primarily consists of members of family Buthidae, is relatively well known (Vachon, 1952; Kovařík, 2006). However, species of the genus *Euscorpius* Thorell from North Africa have not been adequately characterized, even though records from the region date back to more than 100 years. Original reports documented "*E. carpathicus* (L.)" from isolated localities along the North African coast in Tunisia, Libya, and Egypt (Fet et al., 2003). Many *Euscorpius* spp. are known to disperse with humans (Fet et al., 2006), so the legitimacy of these reports has been controversial. Some introduced species, such as *E. italicus* (Herbst) in Yemen and Iraq, are even known to establish reproducing populations in non-native areas (Fet & Kovařík, 2003). Furthermore, some of the African populations of *Euscorpius* are represented by *E. flavicaudis* De Geer, a potential postglacial relict that presumably represents a recent introduction (Gantenbein et al., 2001). As a result, when specimens identified as "*E. carpathicus sicanus* (C.L. Koch)" were reported from coastal regions of North Africa, it was brought into question whether the specimens were introduced from the northern Mediterranean, or if they represented an isolated relict population (Fet et al., 2003).

Based on morphological and molecular characters, *E. carpathicus sicanus* was recently elevated to *E. sicanus* (C.L. Koch), and the degree of intraspecific genetic structure suggested that it might even represent a species complex (Fet et al., 2003). With the type locality from Sicily, and other populations occupying portions of southern Italy, Sardinia, central and southern Greece, Malta, Madeira, and several North African localities, the geographic range of *E. sicanus* is highly fragmented by the Mediterranean Sea. Genetic samples (mitochondrial DNA) of *E. sicanus* were studied from a number of localities in Italy (including Sicily and Sardinia), Greece, and Malta (Fet et al., 2003; Salomone et al., 2007), but no African populations were analyzed.

In 2008, we (P. Stoev & N. Akkari) collected new *Euscorpius* specimens from North Africa that were identified in 2009 as *E. sicanus* (det. V. Fet). The scorpions were collected from Jebel Zaghoun (Fig. 1), a mountain range situated in northeastern Tunisia that reaches an elevation of 1,295 meters at Ras el Gossa. The mountain range is within the Semi-arid bioclimatic zone (Emberger, 1966) characterized by temperate winters and an average annual precipitation of 450–500 mm. Jebel Zaghoun lies in the major structural NE-SW lineament that was active since the Jurassic and is characterized by a predominance of red soils developed on Jurassic limestone. The vegetation near the summit is mostly dominated by *Quercus coccifera* L., the slopes are characterized by *Ceratonia siliqua* L., *Olea europaea* L. and *Pistacia lentiscus* L., and the shrub floor is composed mainly of *Tetraclinis articulata* (Vahl), *Phillyrea angustifolia* L., *Lavandula* sp. and *Thymus capitatus* (L.).

This non-desert habitat suggests that *Euscorpius* from Jebel Zaghoun could potentially represent native populations. We tested this hypothesis by comparing a DNA barcode (mitochondrial *cox1*) from one of the *E. sicanus* specimens (Fig. 2) collected from the Jebel Zaghoun of Tunisia with barcodes obtained from *E. sicanus* from Greece, Malta, and Sardinia, as well as outgroup congeneric species from southern Europe (Fig. 3). We used these data to investigate the matrilineal phylogeny, and to estimate divergence dates between mitochondrial lineages. If *E. sicanus* was recently introduced to North Africa, then we would expect the barcode from the Tunisian sample to be similar to that from Sardinia, Malta, or Greece. Alternatively, if the Tunisian specimen represents a relict population, then we would expect the barcode to be quite different than the *E. sicanus* barcodes from Greece, Sardinia, and Malta. Furthermore, molecular clock

estimates should indicate an ancient (Pre-Pleistocene) divergence between the sample from Tunisia, and those from Greece, Malta, and Sardinia.



Fig. 1. A view of Jebel Zaghouan Mts. in Tunisia where *Euscorpius sicanus* (C.L. Koch) was collected. Photo: N. Akkari.

## Material and Methods

We analyzed 10 sequences obtained at Marshall University (V. Fet; two specimens from Greece) and the Biodiversity Institute of Ontario, University of Guelph, Guelph, Ontario, Canada (G. Blagoev; all other specimens). Label data of the specimens used for DNA analysis are listed below. All sequence data were submitted to GenBank and can be accessed through BOLD (<http://www.boldsystems.org>, Ratnasingham & Hebert, 2007) under project “Scorpions of the Ancient Mediterranean 2 (AMSCO)”. Voucher specimens are in a private collection of V. Fet and in the Biodiversity Institute of Ontario.

**Material Examined:** *Euscorpius sicanus* (C.L. Koch, 1837): GREECE, Thessaly, Mt. Pelion, Visitsa, 39°20'N, 23°08'E, 7 May 2001, leg. V. Fet, VF-0454 (JX414017); Thessaly, Mt. Ossa (Kissavos), Spilia, 39°47'45"N, 22°38'49"E, 9 May 2001, leg. V. Fet, VF-0455 (JX414018); ITALY, Sardinia, S. Niccolo Gerrei, near Grotta Saturru, 39.49816°N, 09.31503°E, 395 m, April 2006, leg. A. v.d. Meijden, VF-0789, AMSCO052-10 (JX133089). MALTA, Buskett Gardens, 35°51'41"N, 14°23'56"E, 17 September 2001, leg. P. Schembri, VF-0792, AMSCO053-10 (HM418288). TUNISIA, Zaghouan Governorate, Jebel Zaghouan Mts., along the trek, 36°22.423'N, 10°06'E to 36°22.924'N, 10°06.789'E, 650-780 m a.s.l., mixed forest, March 2008, leg. P. Stoev & N. Akkari, VF-0793, AMSCO054-10 (HM418289). *Euscorpius carpathicus* (L., 1767): ROMANIA, Caraș-Severin County, Băile Herculane, 44°52'43"N, 22°24'51"E, 4 June 2008 (F. Šťáhlavský), VF-0768, AMSCO044-10 (HM418284). *Euscorpius concinnus* (C.L. Koch, 1837): FRANCE, Alpes-Maritimes, Grasse, 43°40'N, 06°55'E, September 2004, leg. E. Ythier, VF-0782, AMSCO049-10 (HM418287). *Euscorpius hadzii*

Caporiacco, 1950: BULGARIA, Blagoevgrad District, Gorna Breznitsa, 41°45'N, 23°07'E, 27 May 2005, leg. V. Fet & D. Dobrev, VF-0798, AMSCO059-10 (HM880289); MONTENEGRO, Budva District, Visnjevo, 42°17'52"N, 18°46'37"E, sea level, 29 October 2005, leg. F. Franeta, VF-0807, AMSCO066-10 (HM418296). *Euscorpius flavicaudis* (DeGeer, 1787): FRANCE, Vaucluse, Pernes-les-Fontaines, 43°59'55"N, 05°03'35"E, 230 June 2007, leg. V. Fet, VF-0700, AMSCO001-10 (HM418267).

NOTE. Additional specimens of *E. sicanus* (not included in the DNA study) were collected from the same area by us (N.A. and P.S.): 2 juv., NE Tunisia, Zaghouan Governorate, Jebel Zaghouan Mts., surroundings of a small limestone cave 'Gouffre du courant d'air', 36°21.980'N, 10°05.513'E, 561 m a.s.l., *Quercus ilex*, *Pistacia lentiscus*, *Jasminum fruticans*, under stones and leaf litter, 17 March 2008, N. Akkari & P. Stoev leg.



Fig. 2. Dorsal view of the habitus of *Euscorpius sicanus* (C.L. Koch) female collected from Tunisia for which a DNA barcode was sequenced and analyzed in this study. Note a weak darker reticulation pattern on carapace, typical of *E. sicanus*. Photo: P. Stoev and R. Bekchiev.

**Molecular Techniques:** The V.F. lab used a DNeasy Blood & Tissue Kit (Qiagen) to isolate genomic DNA from leg or muscle tissue. A portion of the mitochondrial protein-coding gene cytochrome oxidase subunit I (*cox1*) was then amplified and sequenced using primers Nancy (Simon *et al.*, 1994) and LCO (5' – GGT CAA CAA ATC ATA AAG ATA TTG G – 3') following protocols outlined by Simon *et al.* (1994).

Barcodes (*cox1* sequences) generated at the Canadian Centre for DNA Barcoding, University of Guelph, were obtained using standard protocols for DNA extraction, polymerase chain reaction (PCR) and sequencing (Ivanova *et al.*, 2006, DeWaard *et al.*, 2008). In brief, tissue from a single scorpion leg was used for extraction of genomic DNA using a 96 AcroPrep™ 1 ml filter plate (PALL) with 3.0 µm Glass fiber. DNA was eluted in 40 µl of dH<sub>2</sub>O. Full-length *cox1* barcodes (649 bp) were amplified using two newly designed primer sets (Ivanova, unpublished): ScorpF1\_t1 (5' – TGTAACGACGG CCAGTTTTCTACTAATCAYAAAGAYATTGG – 3') and ScorpR1\_t1 (5' – CAGG AAACAGCTATGACGGRTGTCCAAAAAYCAAAAYAAATG – 3'). All PCR products were sequenced bi-directionally on an ABI3730XL using the primer pair of

M13F and M13R (Messing, 1983). The forward and reverse sequences were used to generate a single consensus sequence using CodonCode Aligner v. 3.0.2 (CodonCode Corporation). Cox1 was chosen because it is commonly used in barcoding and has been demonstrated as highly effective in discriminating among insect (Zhang & Hewitt, 1997; Footitt et al., 2009; Zhou et al., 2009) and arachnid species (e.g. Barrett & Hebert, 2005; Thomas & Hedin, 2008; Wang et al., 2008; Robinson et al., 2009; Graham et al., 2012; Sousa et al., 2012).




Fig. 3. Map depicting locations for Euscorpium Thorell specimens used in this study.

**Phylogenetic analysis and divergence time estimation:** Sequences were aligned using SEQUENCHER v. 4.9 (Gene Codes Corp., Inc., Ann Arbor, MI, USA) and verified by eye. The alignment was then imported into the program MEGA 5 (Tamura et al., 2011) which was used to find a suitable model for nucleotide substitution through the Akaike Information Criterion (Posada, 2008). The program chose the GTR+I+G model, so phylogeny was then estimated via this model and the criterion of Maximum Likelihood (ML) with 1,000 bootstrap replicates, again using MEGA 5.

We also estimated tree topology and divergence dates for the Euscorpium samples in BEAST v. 1.5.3 (Drummond & Rambaut, 2007) using the same substitution model. We applied the Yule tree prior and a mutation rate of 0.007 substitutions/site/million years for cox1 (Gantenbein et al., 2005), and set the mean standard deviation to 0.003 to accommodate a similar rate estimated for 16S (Gantenbein & Largiadèr, 2002). Analyses were conducted for 40 million Markov Chain Monte Carlo generations, sampling every 1,000 generations, and with the first 20% of the generations discarded as burn-in. We used LOGCOMBINER v. 1.6.1 (Drummond & Rambaut, 2007) to combine trees and parameter estimates, and TRACER to examine the estimated sample sizes (ESS) to avoid poor estimates of the parameters (ESS < 200).

## Results

ML and Bayesian analyses produced identical topologies. We chose to present the Bayesian tree with both posterior probabilities and bootstrap support values for each node (Fig. 4, Table 1). A total of 6 out of 9 nodes were supported under BI (PP > 0.9), and 5 nodes were supported by the ML (bootstrap values > 0.75).

Table 1. Molecular clock estimates and support values for nodes presented in Fig. (4).

Node	Age	95% HPD	Posterior Probability	ML Bootstrap (%)
a	25.57	14.32 - 39.56	1	100
b	15.28	9.74 - 22.7	1	100
c	12.65	8.23 - 18.54	0.74	59
d	9.73	5.38 - 15.4	0.98	64
e	8.59	5.52 - 12.25	1	81
f	7.18	4.44 - 10.13	0.66	37
g	5.56	3.29 - 8.14	0.81	38
h	4.67	2.7 - 6.99	1	98
i	3.57	1.72 - 5.67	1	79




Fig. 4. Ultrametric tree estimated in BEAST. Mean divergence times, 95% highest posterior densities (HPD), and support values for nodes (a - i) are presented in Table 1. Dark bars represent variation (95% HPD) for the age estimate of each node.

The tree is rooted with *Euscorpius flavicaudis*, which was estimated to have diverged from the remaining samples sometime between the mid-Oligocene and mid-

Miocene. The next oldest node diversified between the Late Oligocene and Middle Miocene, resulting in two clades: one that was strongly supported by BI but weakly supported by ML consisting of *E. concinnus* and *E. carpathicus*, and another showing sister relationships between *E. hadzii* and *E. sicanus*, which is supported by morphological data (Fet & Soleglad, 2002, 2007; Fet et al., 2003). The *E. concinnus* and *E. carpathicus* were estimated to have diverged in the Middle to Late Miocene. Within the other clade, *E. hadzii* and *E. sicanus* were estimated to have split sometime in the Middle Miocene. Of the *E. sicanus*, the specimen from Malta was most basal and estimated to have diverged from the rest between the Middle and Late Miocene. Of the remaining *E. sicanus*, two specimens from Greece (eastern Thessaly) formed a strongly supported group that was predicted to have diverged from the Late Miocene to early Pleistocene. Although poorly supported, the specimens from Sardinia and Tunisia grouped together in both analyses and were predicted to have diverged from the other *E. sicanus* in the Late Miocene to early Pliocene. The Tunisia specimen was estimated to have diverged from the specimen from Sardinia sometime between the Late Miocene and late Pliocene, with a mean divergence date estimate of 5.56 Ma (Table 1).

## Discussion

In a review of *E. sicanus*, Fet et al. (2003) wrote that “No DNA is available from the northern African enclaves yet; it remains to be seen if these are true relict populations or if they have been introduced via human activity.” The analyses presented herein support the former hypothesis, that North African *E. sicanus* from Tunisia are genetically distinct and represent a relict population. Our sample of *E. sicanus* from Tunisia grouped most closely with a sample from Sardinia. Both Sardinia and Tunisia samples were more distantly related to samples from Greece and Malta. Molecular dating estimated samples from Sardinia and Tunisia to have diverged between the Late Miocene (3.29 Mya) and early Pliocene (8.14 Mya), with a mean estimate near the Mio-Pliocene transition (5.56 Mya), suggesting that the disjunction across the Mediterranean Sea is quite ancient (Fig. 4, Table 1). Intriguingly, this timeframe very closely matches that of a widespread drying and refilling of the Mediterranean Basin in the late Miocene (more precisely the Messinian).

Approximately 5.96 Mya, marine gateways between the Atlantic Ocean and Mediterranean Sea closed due to uplift along the African and Iberian continental margins (Duggen et al., 2003). This resulted in a pervasive desiccation of the Mediterranean Basin known as the ‘Messinian salinity crisis’, which was one of the most dramatic earth history events during the Cenozoic era (Krijgsman, 2002). Evaporation of the Mediterranean Sea is thought to have allowed many terrestrial organisms that were previously isolated by marine waters (e.g. Martín-Piera & Sanmartín, 1999; Sanmartín, 2003; Wilke, 2003), to more easily disperse throughout the region. Tectonic subsidence then allowed Atlantic water to make its way through the Gibraltar Strait at 5.33 Mya. This refilling of the Mediterranean Basin, known as the ‘Zanclean’ or ‘post-Messinian’ flood, then appears to have caused vicariance between terrestrial organisms in North Africa and Europe (Sanmartín, 2003). Such a scenario could account for the genetic divergence between *E. sicanus* from Tunisia and Sardinia. Although we have not studied samples from the Italian mainland (Apennine Peninsula), paleogeographic reconstructions suggest that terrestrial connections occurred between Italy, North Africa, Sicily, Sardinia, and Corsica until the Late Miocene or Pliocene (Rosenbaum & Lister, 2002). Therefore, *E. sicanus* may have dispersed between these regions, which may have been made even easier during the Messinian salinity crisis. Increased longitudinal crustal extension could have then worked synergistically with the refilling of the Mediterranean

basin to effectively sever land connections between our samples from Sardinia and Tunisia, which is concordant with our estimated divergence dates (Fig. 4, Table 1).

If the Zanclean flood was responsible for vicariance in *Euscorpius*, then our rate-calibrated molecular clock was remarkably accurate. Therefore, for similarly distributed taxa (in North Africa and Sardinia) that lack reliable rates, we propose that the Zanclean flood could potentially be used as an incredibly precise geologic calibration. Paleogeographic events like uplift and marine transgressions have commonly been used to date vicariant events, but these events happen gradually and the actual timing of the reduction in gene flow cannot be pinpointed. However, the Zanclean Flood is thought to have filled the Mediterranean in 2 months to 2 years (Garcie-Castellanos et al., 2009), and as similarly proposed for river capture and reversals with freshwater-limited organisms (Waters et al., 2007), the event could potentially be used as a ‘sharp’ vicariant event, allowing for more precise calibrations. Other authors have already used this sharp vicariant event to calibrate molecular clocks for organisms in the eastern Mediterranean, as it is thought to have isolated populations on Crete and Cyprus (e.g. Beerli et al., 1996; Gantenbein & Keightley, 2004; Lymberakis et al., 2007; Akin et al., 2010; Kornilios et al., 2012). As far as we are aware, however, this method has not yet been employed for organisms from Tunisia and Sardinia.

The placement of our sample from Malta as the most basal lineage within *E. sicanus* is curious. Based on mtDNA data from 16S (Fet et al., 2003), the same specimen is most closely related to samples from Nebrodi, Sicily, which is the type locality for *E. sicanus*. Although Malta is closer to Sicily than Sardinia and Tunisia, this relationship is somewhat surprising when considering earth history. As mentioned above, land connections occurred between Sardinia, Corsica, Sicily, and Tunisia until the Late Miocene to early Pliocene. Although an underwater ridgeline connects the Maltese Islands with Sicily and Tunisia, paleogeographic reconstructions suggest that the archipelago may have remained insular for at least several million years longer (Rosenbaum & Lister, 2002). Therefore, the Maltese Islands could have been colonized by mainland populations of *E. sicanus* that dispersed to the islands prior to the Messinian salinity crisis. Alternatively, *E. sicanus* could have colonized the island of Malta and dispersed to the mainland, probably Sicily, where it may now occur in sympatry with other lineages (represented by our sample in Sardinia) that diverged during the Zanclean Flood. Additional sampling along the along the Apennine Peninsula, Sicily, and the remaining Maltese Islands would be needed to address this hypothesis.

Whatever the mechanism, DNA barcodes imply that North African populations of *E. sicanus* were probably not recently introduced and instead represent an ancient and isolated natural population. If *E. sicanus* had recently colonized the area via human introduction, then the *cox1* barcode should have been similar to those from *E. sicanus* collected in Malta, Greece, or Sardinia, from which the Tunisian population would have most likely been founded. However, we recognize that our sampling is limited, especially in Italy, and that additional cryptic lineages could occur within the species, so recent colonization of North Africa should not be completely ruled out. Furthermore, the age of the intraspecific lineages recovered in *E. sicanus* (some with estimates in the Miocene) suggest that the species might actually represent a cryptic species complex, calling attention to the need for a rigorous and comprehensive assessment of the genus *Euscorpius*. To date, most systematic studies of *Euscorpius* have focused on western Mediterranean and central European species (Gantenbein et al., 2000, 2001; Fet et al., 2003; Salomone et al., 2007). However, recent work has revealed that *Euscorpius* is most diverse in the poorly studied eastern Mediterranean, especially the Balkans, the Aegean region, and Anatolia (Fet et al., in progress). Finally, our analyses provide yet another

example of how DNA barcodes can be used for more than just identifying and discovering species (Hebert et al., 2003; Stoeckle, 2003), and that ‘sharp’ vicariant events like the Zanclean Flood may be useful for fine-tuning molecular clocks.

### Acknowledgments

The DNA barcoding conducted for this project was performed at the Canadian Centre of DNA Barcoding, Biodiversity Institute of Ontario, University of Guelph, with administrative support from P.D.N. Hebert, and funded by the Government of Canada through Genome Canada and the Ontario Genomics Institute (2008-OGI-ICI-03). We thank the Lead DNA Scientist Natalia Ivanova, Biodiversity Institute of Canada, University of Guelph, Guelph, Ontario, for her expert help and guidance. For collection and donation of *Euscorpius* specimens we are grateful to Filip Franeta, Arje van der Meijden, Patrick Schembri, František Šťáhlavský, and Eric Ythier. P. Stoev and N. Akkari's field trip in Tunisia was supported by the Field Museum Collection Fund, with the logistic help of Petra Sierwald. Alexi Popov, Christo Deltshev, Dobrin Dobrev and Ivan Pandourski provided expert help and transportation for Victor, Galina, Elizabeth, and Simon Fet during their 2005 field trips to collect *E. hadzii* in Bulgaria. The *E. flavicaudis* were collected on the property of Annette and Bernard Janin, who kindly hosted Victor and Galina Fet in 2007 at the wonderful village of Pernes-les-Fontaines in Provence. Travel of Victor and Galina Fet to Bulgaria in 2005 was supported by a Fulbright Foundation grant to V.F. Travel of Victor, Galina and Elizabeth Fet to the University of Guelph in 2009 was supported by Marshall University. Rostislav Bekchiev kindly assisted in photographing *E. sicanus*. We thank Benjamin Gantenbein-Ritter for important comments on the manuscript.

### References

- Akın, Ç., Bilgin, C., Beerli, P., Westaway, R., Ohst, T., Litvinchuk, S.N., Uzzell, T., Bilgin, M., Hotz, H., Guex, G.-D. & Plötner, J. 2010. Phylogeographic patterns of genetic diversity in eastern Mediterranean water frogs were determined by geological processes and climate change in the Late Cenozoic. *Journal of Biogeography*, 37: 2111–2124.
- Barrett, R.D.H. & Hebert, P.D.N. 2005. Identifying spiders through DNA barcodes. *Canadian Journal of Zoology*, 83: 481–491.
- Beerli, P., Hotz, H., Uzzell, T. 1996. Geological dated sea barriers calibrate a protein clock for Aegean water frogs. *Evolution*, 50: 1676–1687.
- DeWaard, J.R., Ivanova, N.V., Hajibabaei, M. & Hebert, P.D.N. 2008. Assembling DNA barcodes: analytical protocols. In: Martin, C.C. (ed), *Methods in Molecular Biology*. Volume 410: *Environmental Genomics*. Pp. 275–283. Humana Press, Totowa, New Jersey.
- Drummond, A.J. & Rambaut, A. 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology*, 7: 214.
- Duggen, S., Hoernle, K., van den Bogaard, P., Rupke, L. & Phipps Morgan, J. 2003. Deep roots of the Messinian salinity crisis. *Nature*, 422: 602–606.
- Emberger, L. 1966. Une classification biogéographique des climats. *Recherches et Travaux des Laboratoires de Géologie, Botanique et Zoologie* vol. 7. 43 pp. Faculté des Sciences Montpellier, France.
- Fet, V., Gantenbein, B., Karataş, Ay. & Karataş, A. 2006. An extremely low genetic divergence across the range of *Euscorpius italicus* (Scorpiones: Euscorpiidae). *Journal of Arachnology*, 34(1): 248–253.
- Fet, V. & Kovařík, F. 2003. A record of *Euscorpius* (Polytrichobothrius) *italicus* (Herbst, 1800) (Scorpiones: Euscorpiidae) in Iraq. *Acta Societatis Zoologicae Bohemicae*, 67: 179–181.

- Fet, V. & Soleglad, M.E. 2002. Morphology analysis supports presence of more than one species in the “*Euscorpius carpathicus*” complex (Scorpiones: Euscorpiidae). *Euscorpius*, 3: 1–51.
- Fet, V. & Soleglad, M.E. 2007. Fauna and zoogeography of scorpions (Arachnida: Scorpiones) in Bulgaria. In: Fet, V. & A. Popov (eds.), *Biogeography and Ecology of Bulgaria*. pp. 405–422. Springer.
- Fet, V., Soleglad, M.E., Gantenbein, B., Vignoli, V., Salomone, N., Fet, E.V. & Schembri, P.J. 2003. New molecular and morphological data on the “*Euscorpius carpathicus*” species complex (Scorpiones: Euscorpiidae) from Italy, Malta, and Greece justify the elevation of *E. c. sicanus* (C. L. Koch, 1837) to the species level. *Revue suisse de Zoologie*, 110(2): 355–379.
- Footitt, R.G., Maw, H.E.L., Havill, N.P., Ahen, R.G. & Montgomery, M.E. 2009. DNA barcodes to identify species and explore diversity in the Adelgidae (Insecta: Hemiptera: Aphidoidea). *Molecular Ecology Resources*, 9(1): 188–195.
- Gantenbein, B., Fet, V., Barker, M. & Scholl, A. 2000. Nuclear and mitochondrial markers reveal the existence of two parapatric scorpion species in the Alps: *Euscorpius germanus* (C. L. Koch, 1837) and *E. alpha* Caporiacco, 1950, stat. nov. (Scorpiones, Euscorpiidae). *Revue suisse de Zoologie*, 107(4): 843–869.
- Gantenbein, B., Fet, V., Gantenbein-Ritter, I.A. & Balloux, F. 2005. Evidence for recombination in scorpion mitochondrial DNA (Scorpiones: Buthidae). *Proceedings of the Royal Society B-Biological Sciences*, 272: 697–704.
- Gantenbein, B. & Keightley, P.D. 2004. Rates of molecular evolution in nuclear genes of East Mediterranean scorpions. *Evolution*, 58: 2486–2497.
- Gantenbein, B. & Largiadèr, C.R. 2002. *Mesobuthus gibbosus* (Scorpiones: Buthidae) on the island of Rhodes – hybridization between Ulysses’ stowaways and native scorpions? *Molecular Ecology*, 11(5): 925–938.
- Gantenbein, B., Soleglad, M.E. & Fet, V. 2001. *Euscorpius balearicus* Caporiacco, 1950, stat. nov. (Scorpiones: Euscorpiidae): molecular (allozymes and mtDNA) and morphological evidence for an endemic Balearic Islands species. *Organisms, Diversity & Evolution*, 1(4): 301–320.
- Garcie-Castellanos, D., Estrada, F., Jiménez-Munt, I., Groini, C., Fernández, M., Vergés J. & de Vincete, R. 2009. Catastrophic flood of the Mediterranean after the Messinian salinity crisis. *Nature*, 462: 778–781.
- Graham, M.R., Oláh-Hemmings, V. & Fet, V. 2012. Phylogeography of co-distributed dune scorpions identifies the Amu Darya River as a long-standing component of Central Asian biogeography. *Zoology in the Middle East*, 55: 95–110.
- Hebert, P.D.N., Cywinska, A., Ball, S.L. & deWaard, J.R. 2003. Biological identifications through DNA barcodes. *Proceedings of the Royal Society B-Biological Sciences*, 270: 313–321.
- Ivanova, N.V., DeWaard, J.R. & Hebert, P.D.N. 2006. An inexpensive, automation-friendly protocol for recovering high-quality DNA. *Molecular Ecology Notes*, 6: 998–1002.
- Kornilios, P., Ilgaz, Ç., Kumlutaş, Y., Lymberakis, P., Moravec, J., Sindaco, R., Rastegar-Pouyani, N., Afroosheh, M., Giokas, S., Fraguedakis-Tsolis, S. & Chondropoulos, B. 2012. Neogene climatic oscillations shape the biogeography and evolutionary history of the Eurasian blindsnake. *Molecular Phylogenetics and Evolution*, 62(3): 856–873.
- Kovářík, F. 2006. Review of Tunisian species of the genus *Buthus* with descriptions of two new species and a discussion of Ehrenberg’s types (Scorpiones: Buthidae). *Euscorpius*, 34: 1–16.
- Krijgsman, W. 2002. The Mediterranean: Mare Nostrum of Earth sciences. *Earth and Planetary Science Letters*, 205: 1–12.
- Lymberakis, P., Pouloukakis, N., Manthalou, G., Tsigenopoulos, C.S., Magoulas, A. & Mylonas, M. 2007. Mitochondrial phylogeography of *Rana* (Pelophylax) populations in the Eastern Mediterranean region. *Molecular Phylogenetics and Evolution*, 44: 115–125.

- Martín-Piera, F. & Sanmartín, I. 1999. Biogeografía de áreas y biogeografía de artrópodos Holárticos y Mediterráneos. In: *Evolution and Phylogeny of Arthropoda*. pp. 535–560. Boletín SEA, 26. Sociedad Entomológica Aragonesa, Zaragoza, Spain.
- Messing, J. 1983. New M13 vector for cloning. *Methods in Enzymology*, 101: 20–78.
- Posada, D. 2008. jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution*, 25: 1253–1256.
- Ratnasingham, S. & Hebert, P.D.N. 2007. BOLD: The Barcode of Life Data System ([www.barcodinglife.org](http://www.barcodinglife.org)). *Molecular Ecology Notes*, 7: 355–364. DOI: 10.1111/j.1471-8286.2006.01678.x
- Robinson, E.A., Blagoev, G.A., Hebert, P.D.N. & Adamowicz, S.J. 2009. Prospects for using DNA barcoding to identify spiders in species-rich genera. *ZooKeys*, 16: 27–46. doi: 10.3897/zookeys.16.239
- Rosenbaum, G. & Lister, G.S. 2002. Reconstruction of the evolution of the Alpine-Himalayan orogen - an introduction. *Journal of the Virtual Explorer*, 8: 1–2.
- Salomone, N., Vignoli, V., Frati, F., & F. Bernini. 2007. Species boundaries and phylogeography of the “*Euscorpius carpathicus* complex” (Scorpiones: Euscorpiidae) in Italy. *Molecular Phylogenetics and Evolution*, 43: 502–514.
- Sanmartín, I. 2003. Dispersal vs. vicariance in the Mediterranean: historical biogeography of the Palearctic Pachydeminæ (Coleoptera, Scarabaeoidea). *Journal of Biogeography*, 30: 1883–1897.
- Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H. & Flook, P. 1994. Evolution, weighting, and phylogenetic Utility of mitochondrial gene dequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America*, 87(6): 651–701.
- Sousa, P., Harris, D.J., Froufe, E. & van der Meijden, A. 2012. Phylogeographic patterns of *Buthus* scorpions (Scorpiones: Buthidae) in the Maghreb and South-Western Europe based on CO1 mtDNA sequences. *Journal of Zoology*, in press (first published online: 16 May 2012).
- Stoeckle, M. 2003. Taxonomy, DNA and the bar code of life. *BioScience*, 53: 2–3.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. 2011. MEGA 5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28: 2731–2739.
- Thomas, S.M. & Hedin, M. 2008. Multigenic phylogeographic divergence in the paleoendemic southern Appalachian opilionid *Fumontana deprehendor* Shear (Opiliones, Laniatores, Triaenonychidae). *Molecular Phylogenetics and Evolution*, 46: 645–658.
- Vachon, M. 1952. *Etudes sur les Scorpions*. 482 pp. Institut Pasteur d’Algérie, Alger.
- Wang, Q., Li, S., Wang, R. & Paquin, P. 2008. Phylogeographic analysis of Pimoidae (Arachnida: Araneae) inferred from mitochondrial cytochrome c oxidase subunit I and nuclear 28S rRNA gene regions. *Journal of Zoological Systematics and Evolutionary Research*, 46: 96–104.
- Waters, J.M., Rowe, D.L., Apte, S., King, T.M., Wallis, G.P., Anderson, L., Norris, R.J., Craw, D. & BurrIDGE, C.P. 2007. Geological dates and molecular rates: rapid divergence of rivers and their biotas. *Systematic Biology*, 56: 271–282.
- Wilke, T. 2003. *Salenthydrobia* gen. nov. (Rissooidea: Hydrobiidae): a potential relict of the Messinian salinity crisis. *Zoological Journal of the Linnean Society*, 137: 319–336.
- Zhang, D.-X. & Hewitt, G.M. 1997. Assessment of the universality and utility of a set of conserved mitochondrial primers in insects. *Insect Molecular Biology*, 6: 143–150.
- Zhou, X., Adamowicz, S.J., Jacobus, L.M., DeWalt, R.E. & Hebert, P.D.N. 2009. Towards a comprehensive barcode library for arctic life - Ephemeroptera, Plecoptera, and Trichoptera of Churchill, Manitoba, Canada. *Frontiers in Zoology*, 6: 1–9.